



<b>PRODUCE:</b>	<b>TOMATO</b>
-----------------	---------------

<b>TYPE</b>	<b>Truss</b>	<b>VARIETY</b>	<b>Various</b>
<b>CLASS</b>	<b>One</b>	<b>NOTES</b>	

GENERAL APPEARANCE CRITERIA	
<b>Colour</b>	With receival colour as pre-ordered; generally with first tomato fully red (stage 5-6 on QDPI Chart), last tomato > % red colour (stage 2 on QDPI Chart); nil with mottled appearance.
<b>Visual Appearance</b>	Minimum 4 tomatoes attached to a central branch; bright green calyxes and stems; bright bloom on fruit skins; pale seeds in clear, reddish gel.
<b>Sensory</b>	Thin, smooth skin; firm flesh with rich tomato flavour; no foreign odours or tastes.
<b>Shape</b>	Uniformly round; nil kidney, egg or dumb-bell shaped, or with obvious raised ridges, grossly uneven segments or otherwise deformed
<b>Size</b>	Minimum 4 tomatoes per truss, (no more than 1 truss in tray with a minimum 3 tomatoes); tomato diameter 55 - 75mm; as per pre-ordered requirements.
<b>Maturity</b>	Firm, vine ripened fruit (no gassed fruit); no fully green (underripe) or soft (overripe) fruit.
MAJOR DEFECTS	
<b>Insects</b>	With obvious live insects (eg insect larvae)
<b>Diseases</b>	With fungal or bacterial rots (eg bacterial soft rot, Alternaria rot, grey mould, sour rot). With blotchy or mottled appearance due to viral infection.
<b>Physical/Pest Damage</b>	With cuts, holes, splits or insect damage that breaks the skin. With soft, dark areas due to severe bruising.
<b>Physiological Disorder</b>	With softening or rot developing at the blossom end (calcium deficiency). With grey or brown areas in the internal flesh (grey wall / vascular browning). With circular or star shaped cracking around the stem attachment or blossom scar.
<b>Temperature Injury</b>	With softened flesh, pitted skin, or failure to develop normal colour (chilling injury). With bleached, flattened areas on the fruit skin (sunburn). With soft skin or wrinkled appearance (dehydration) With blotchy, orange toned skin (heat damage).
MINOR DEFECTS	
<b>Skin Marks/Blemishes</b>	With light scattered blemish, eg. rub marks, affecting in aggregate > 1 sq cm. With brown marks or flesh discolouration affecting in aggregate > 0.5 sq cm. With 'zipper' lines > half the fruit length and >1 mm width.
<b>Physical/Pest Damage</b>	With minor bruises or impact damage eg skin slightly dark but still firm, affecting >2 sq cm.
<b>Physiological Disorder</b>	With persistent green shoulders affecting >2 sq cm.
CONSIGNMENT CRITERIA	
<b>Tolerance Per Consignment</b>	Total minor defects (within allowance limit) to be < 2 defects per item Total minor defects (outside allowance limit) must not exceed 10% of consignment. Total major defects must not exceed 2 % of consignment. Combined Total not to exceed 10%.
<b>Packaging &amp; Labelling</b>	Packaging manufactured from new food grade materials or sanitised returnable crates. All labelling must meet the current legislative requirements. Labelling to identify grower's name/brand (plus growers name/code if via a packhouse), address, contents, class, size and/or minimum net weight. Produce to identify Country of Origin (eg. Produce of Australia) on outer container..
<b>Shelf Life</b>	Produce must provide not less than 14 days clear shelf life from date of receival.
<b>Receival Conditions</b>	Compliance with Quarantine Treatments (if required) for Interstate Consignment. Stacked onto a stabilised pallet as pre-ordered. Refrigerated van with air bag suspension, unless otherwise approved. Pulp Temperature 12 - 15° C.
<b>Chemical &amp; Containment Residues</b>	All chemicals used pre/postharvest must be registered and approved for use in accordance with the requirements of the APVMA regulatory system. Residues, Contaminants and Heavy Metals to comply with the FSA NZ Food Standards Code MRL's and ML's.
<b>Food Safety Requirements</b>	Produce is to be grown and packed under a HACCP based food safety program that is subject to an annual third-party audit. A copy of current certification to be forwarded to receiver.