



<b>PRODUCE:</b>	<b>POTATO</b>
-----------------	---------------

<b>TYPE</b>	<b>Washed, Pre Pack (6kg &amp; 10kg)</b>	<b>VARIETY</b>	<b>Pontiac</b>
<b>CLASS</b>	<b>Composite</b>	<b>NOTES</b>	

GENERAL APPEARANCE CRITERIA	
<b>Colour</b>	With deep pink to red skin; white-cream flesh.
<b>Visual Appearance</b>	Medium depth eyes; slightly rough skin; free from foreign matter. Tubers to be free from excessive moisture, not wet.
<b>Sensory</b>	Firm when tested with finger pressure, not soft or spongy; free from foreign tastes or odours.
<b>Shape</b>	Approximately round to oblong.
<b>Size</b>	100 - 400g with 50% in range 130 - 340g.
MAJOR DEFECTS	
<b>Insects</b>	With obvious live insects (eg potato moth, potato wireworm).
<b>Diseases</b>	With fungal or bacterial rots (eg pink rot, dry rot, gangrene, late blight or bacterial wilt).
<b>Physical/Pest Damage</b>	With evidence of insect, rodent, or nematode damage. With cuts, splits, cracks or other wounds >40mm long, and/or >4mm wide and/or >4mm deep With swollen lenticels (blistered/watery) >2mm in diameter affecting >7sq cm. With severely malformed or root constricted tubers, or tubers with advanced secondary growth. With deep, dark bruising and softened flesh. With green areas on the skin or an overall green tinge (exposed to light, toxic compounds may form).
<b>Physiological Disorders</b>	With sprouting from the eyelets (overstored tubers). With severe brown spotting of the flesh (brown fleck) or hollow cavities inside the tubers (hollow heart). With dark pink or black areas at the centre of the tuber (black heart, due to low oxygen).
<b>Temperature Injury</b>	With reddish brown blotches or soft, water-soaked areas in the flesh (freezing injury). With obvious shrivelling and softened, spongy flesh (dehydration).
MINOR DEFECTS	
<b>Physical/Pest Damage</b>	With cuts or splits <40mm long, <4mm wide and <4mm deep (not removable by one pass peeling). With superficial bruises, scuffs, pressure or rub marks affecting in total >4sq cm and 4mm deep.
<b>Skin Marks/Blemishes</b>	With moderate development of a silvery sheen, brown markings or skin cracking (crazing) affecting >6sq cm. With raised or sunken corky growths on the potato skin (powdery scab, common scab) > 5 sq cm With moderate black speckling affecting in aggregate >2 sq cm of surface area.
<b>Physiological Disorder</b>	With enlarged lenticels (>2mm diameter) affecting >7sq cm. With healed growth cracks or splits >5mm wide or >5mm deep.
CONSIGNMENT CRITERIA	
<b>Tolerance Per Consignment</b>	Total minor defects (within allowance limit) to be < 2 defects per item Total minor defects (outside allowance limit) must not exceed 10% of consignment. Total major defects must not exceed 2% of consignment. Combined Total not to exceed 10%
<b>Packaging &amp; Labelling</b>	Packaging manufactured from new food grade materials or sanitised returnable crates. All labelling must meet the current legislative requirements. Labelling to identify grower's name/brand (plus growers name/code if via a packhouse), address, contents, class, size and/or minimum net weight. Produce to identify Country of Origin (eg. Produce of Australia) on outer container.
<b>Shelf Life</b>	Produce must provide not less than 14 days clear shelf life from date of receipt.
<b>Receival Conditions</b>	Compliance with Quarantine Treatments (if required) for Interstate Consignment. Stacked onto a stabilised pallet as pre-ordered. Refrigerated van with air bag suspension, unless otherwise approved. Pulp Temperature 12 - 20°C
<b>Chemical &amp; Containment Residues</b>	All chemicals used pre/postharvest must be registered and approved for use in accordance with the requirements of the APVMA regulatory system. Residues, Contaminants and Heavy Metals to comply with the FSANZ Food Standards Code MRL's and ML's.
<b>Food Safety Requirements</b>	Produce is to be grown and packed under a HACCP based food safety program that is subject to an annual third-party audit. A copy of current certification to be forwarded to receiver.